

**COMMENSALS ON NESTING HAWKSBILL TURTLES (*ERETMOCHELYS IMBRICATA*), MILMAN ISLAND, NORTHERN GREAT BARRIER REEF, AUSTRALIA.** *Memoirs of the Queensland Museum* 49(2): 674. 2004. - Sea turtles carry a large array of commensals (Frazier et al., 1985; et al., 1991; et al., 1992; Hollenberg & Norris, 1977; Monroe & Limpus, 1979) that are often overlooked. Witzell (1983) noted the incomplete commensal data set for hawksbill turtles (*Eretmochelys imbricata*). Studies of hawksbill turtle nesting patterns and behaviours at Milman Island (Dobbs et al., 1999; Loop, 1996; Loop et al., 1995) allowed collection of commensals during summers January 1991- February 1995.

Commensals were found on 88% (n=1225) of 1392 nesting hawksbill turtles. The most frequent commensal was the barnacle *Chelonibia caretta*, found on the carapace, plastron or head of 1123 turtles (81%); egg-bearing *C. caretta* were observed December 1994 - February 1995. *Balanus* sp. (carapace) and *Platylepas* sp. (skin) were other barnacles randomly noted on nesting hawksbills. The burrowing barnacle *Tubicinella cheloniae* was found in the carapace of 30% of nesting turtles (n=422). Barnacles were encountered at all locations on the carapace. No consistent barnacle attachment pattern was noted. On some turtles, *Tubicinella cheloniae* covered up to 75% of the carapace, while some burrowed through the edge of marginal scutes.

Filamentous algae occurred on 47% (n=649) of nesting turtles. *Gelidiopsis intricata*, *Gelidiella pannosa* and *Polysiphonia* sp. were identified. Nongeniculate corallinaceae (encrusting algae) was found on the carapace or on attached barnacles of 18 turtles (1.3%). Algae was encountered everywhere on the carapace and predominantly under the trailing edge of the last vertebral and costal scutes.

*Ozobranchus* sp. leeches occurred around the cloaca, attached to the underside of front flippers, around a crater wound or on the face of 25 hawksbills (1.8%). The only period during which leech eggs were recorded was January - April 1993. The eggs were on the posterior ventral end of the carapace and on the plastron of nesting turtles. Marine worms (Cirratulidae, Eunicidae, Nereididae) occurred on 18 turtles (1.3%). Other items on the carapaces included a vermetid gastropod (2 turtles), marine crabs (4 turtles), Ophiurid brittlestars (2 turtles), half of an oyster shell (1 turtle), mud (2 turtles) and amphipods.

Commensals on most hawksbills at Milman Island were similar to those noted by Limpus et al., (1983) who reported 100% of their nesting conspecifics had commensal barnacles. Diamond (1976), Pritchard (1969) and Ross (1981) found most hawksbill turtles nesting at Cousin Island, Seychelles, Guyana and Oman, respectively, had large barnacles on their carapaces. Vaughan (1981) reported commensals from 55% of Solomon Island hawksbills. All species of barnacle recorded on hawksbills herein had been reported by Monroe & Limpus (1979).

Hawksbill turtles range over a variety of habitats and may migrate long distances between foraging and nesting areas (Miller et al., 1998; Parmenter, 1983; Vaughan & Spring, 1980). Their preferred foraging habitats commonly include rocky areas, coral reefs, lagoons, bays and estuaries in the circumtropical Atlantic, Indian and Pacific Oceans (Witzell 1983). Terrigenous material, found here on 2 turtles and on 1 at Campbell Island (Limpus et al., 1983) indicates migration from inshore waters where mud may have been encountered whilst foraging. Coral reefs that surround most hawksbill nesting beaches (Witzell, 1983) do not contain mud.

An in-depth analysis of sea turtle commensals, their biology and distribution should provide new insights on habitats in which they have lived. Careful collection and identification of organisms living in association with the

epibiotic communities require further attention. Sadly specimens collected during this study were lost during a flood.

#### Acknowledgements

Identifications of epibionts were conducted by D. Blair (parasites), J. Collins (barnacles) and I. Price (algae), James Cook University, and P. Arnold (worms), Museum of Tropical Queensland. Funding was provided from the Queensland EPA, Jordan Institute for International Awareness, Research Foundation, and Study Abroad Office at Texas A&M University, Institute for International Education, International Women's Fishing Association, and Help Endangered Animal Ridley Turtles (HEART).

#### Literature Cited

- DIAMOND, A.W. 1976. Breeding biology and conservation of hawksbill turtles, *Eretmochelys imbricata* L., on Cousin Island, Seychelles. *Biological Conservation* 9: 199-215.
- DOBBS, K.A., MILLER, J.D., LIMPUS, C.J. & LANDRY, A.M., JR. 1999. Hawksbill turtle, *Eretmochelys imbricata*, nesting at Milman Island, northern Great Barrier Reef, Australia. *Chelonian Conservation and Biology* 3(2): 344-361.
- FRAZIER, J.G., GOODBODY, I. & RUCKDESCHER, C.A. 1991. Epizoon communities on marine turtles: II. tunicates. *Bulletin of Marine Science* 48: 763-765.
- FRAZIER, J., MARGARITOU, D., MULDOON, K., POTTER, C.W., ROSEWATER, J., RUCKDESCHER, C. & SALAS, S. 1985. Epizoon communities on marine turtles I. bivalve and gastropod molluscs. *Marine Ecology* 6: 127-140.
- FRAZIER, J., WINSTON, J.E. & RUCKDESCHER, C.A. 1992. Epizoon communities on marine turtles: III. bryozoa. *Bulletin of Marine Science* 51: 1-8.
- HOLLENBERG, G.J. & NORRIS, J.N. 1977. The red algae *Polysiphonia* (Rhodomelaceae) in the northern Gulf of California. *Smithsonian Contributions in Marine Science*, 1.
- LIMPUS, C.J., MILLER, J.D., BAKER, V. & MCLACHLAN, E. 1983. The hawksbill turtle, *Eretmochelys imbricata* (L.), in North-eastern Australia: the Campbell Island rookery. *Australian Wildlife Research* 10: 185-197.
- LOOP, K.A. 1996. Hawksbill turtle (*Eretmochelys imbricata*) nesting at Milman Island, Queensland, Australia. Unpubl. PhD dissertation, Texas A&M University, Department of Wildlife and Fisheries Sciences, College Station, Texas USA.
- LOOP, K.A., MILLER, J.D. & LIMPUS, C.J. 1995. Nesting by the hawksbill turtle (*Eretmochelys imbricata*) on Milman Island, Northern Great Barrier Reef, Australia. *Wildlife Research* 22: 241-252.
- MILLER, J.D., DOBBS, K.A., MATTOCKS, N., LIMPUS, C.J. & LANDRY, A.M., JR. 1998. Long distance migrations by the hawksbill turtle (*Eretmochelys imbricata*) from Eastern Australia. *Wildlife Research* 25: 89-95.
- MONROE, R. & LIMPUS, C.J. 1979. Barnacles on turtles in Queensland waters with descriptions of three new species. *Memoirs of the Queensland Museum* 19: 197-223.
- PARMENTER, C.J. 1983. Reproductive migration in the hawksbill turtle (*Eretmochelys imbricata*). *Copeia* 1983: 271-3.
- PRITCHARD, P. 1969. Sea turtles of the Guianas. *Bulletin of the Florida State Museum* 13: 85-140.
- ROSS, J.P. 1981. Hawksbill turtle *Eretmochelys imbricata* nesting in the Sultanate of Oman. *Biological Conservation* 19: 99-106.
- VAUGHAN, P. 1981. Marine turtles: a review of their status and management in the Solomon Islands. (Ministry of Natural Resources, Fisheries Division: Honiara, Solomon Islands).
- VAUGHAN, P. & SPRING, S. 1980. Long distance hawksbill recovery. *Marine Turtle Newsletter* 16: 6-7.
- WITZELL, W.N. 1983. Synopsis of biological data on the hawksbill turtle, *Eretmochelys imbricata* (Linnaeus, 1766). *FAO Fisheries Synopsis* 137.1.
- K.A. Dobbs\* & A.M. Landry Jr, *Institute of Marine Life Sciences, Texas A&M University, 4700 Ave U., Bldg 303, Galveston, TX 77551 USA; J.D. Miller, American University in Cairo, PO Box 2511, 113 Sharia Kasr El Aini, Cairo, Egypt; \* present address: Great Barrier Reef Marine Park Authority, PO Box 1379, Townsville 4810, Australia (e-mail: kirstind@gbmpa.gov.au); 16 May 2003.*